



**GLUTEN
INTOLERANCE
GROUP**



Dermatitis Herpetiformis

Updated June 2019

What you need to know about dermatitis herpetiformis

Dermatitis Herpetiformis (DH) is a form of celiac disease. It is a skin condition characterized by groups of itchy blisters. The ingestion of gluten (a protein contained in wheat, rye and barley) triggers an immune system response that deposits IgA antibodies under the top layer of skin. IgA antibodies are present in affected as well as unaffected skin. If you have DH and do not follow a gluten-free diet, you may develop the intestinal damage which is characteristic of celiac disease. In DH, the primary lesion is on the skin rather than the small intestine. The degree of damage to the small intestine is often less severe or more patchy than in those with celiac disease who do not have DH. DH is more common in men than in women, affects approximately 10-15% of people with celiac disease, and is typically diagnosed in adulthood. DH is permanent and symptoms/damage will occur if gluten is consumed.

Symptoms

The IgA deposits result in eruptions of a painfully itchy rash and may progress to red, raised patches of skin that develop into small, watery blisters. The itching and burning of the eruptions are severe and the urge to scratch them is intense. Scratching will further irritate the eruptions. Eruptions commonly occur on pressure points - around the elbows, the front of the knees, the buttocks, back, shoulders, face, and scalp, but can occur anywhere. Eruptions are usually bilateral, occurring on both sides of the body. Obvious gastrointestinal symptoms in DH are uncommon.

Diagnosis

If the eruptions appear to be DH, your dermatologist will take a small biopsy of unaffected skin, next to an eruption. The presence of IgA deposits confirms a diagnosis of DH. An endoscopy is not necessary for diagnosing DH. Only about half of individuals with DH test positive for celiac disease using standard CD blood screening tests. Dermatitis herpetiformis is an inherited autoimmune digestive disease and confirmation of DH will help future generations be aware of the risk within the family.

Treatment

Strictly following a gluten-free diet for life is the only complete treatment. This involves the elimination of wheat, rye, barley, and foods made from these grains

Questions to ask your doctor:

Should I take medication for this disease?

How long will I need to take this medicine and how will I know when to stop taking it?

What are the side effects of these medicines?

How often do I need to get my blood drawn to monitor this medicine's effect on my body?

What else can trigger DH?

Should I take nutritional supplements?

Could I have associated food intolerances?

Where can I have a bone-density study?

How can I find out about the diet?

How often should I follow up with the doctor?

(or their derivatives) from the diet. It may take two or more years for the IgA deposits under the skin to completely clear. A medication called Dapsone may also be prescribed. You will need to follow up with your doctor on a regular basis if you use this drug. Dapsone allows the eruptions to heal, but does not cure DH. Discuss the potential side effects of this drug with your doctor before starting it. Your goal should be to take as little as possible for as short a time as required to allow the diet to control the DH.

The Diet

The gluten-free (GF) diet is a lifelong commitment and should not be started before being properly diagnosed with DH. Starting the diet without complete testing is not recommended and makes later diagnosis difficult. Tests to confirm DH could be negative if a person were on a GF diet for a period of time. For a valid diagnosis, gluten would need to be reintroduced for at least several weeks before testing.

Prognosis

Excellent, if you stay on the gluten-free diet. The severity and frequency of eruptions will decrease as you continue with the diet. Iodine and sun may trigger eruptions in some people. However, iodine is an essential nutrient and should not be removed from the diet without a physician's supervision.

Reference

Salmi, T T. (2019). Dermatitis Herpetiformis. *Clinical and Experimental Dermatology*. <https://onlinelibrary.wiley.com/doi/abs/10.1111/ced.13992>. Accessed 6/12/19.

This document may be reproduced for educational purposes only.

Other helpful information is available at www.GLUTEN.org.

Advances in celiac disease are fast-paced. If this document is more than 2 years old, please visit our website for updated documents.

This information should not be used to diagnose or treat gluten-related disorders or other medical conditions. For questions about these conditions consult your healthcare team when considering this information.

Please consider your local GIG support group as another resource.

Gluten Intolerance Group (GIG)
31214 – 124th Ave. S.E.
Auburn, WA 98092-3667

Phone: 253-833-6655
Fax: 253-833-6675

www.GLUTEN.org
customerservice@GLUTEN.org

The Mission of the Gluten Intolerance Group is to empower the gluten-free community through consumer support, advocacy, and education.

To make a donation or become a volunteer to GIG, visit our website or call the office at 253-833-6655.